

T113
+Y12
3253

YALE UNIVERSITY LIBRARY



3 9002 06679 2079

CONVERSION OF PTEROYL MONOGLUTAMATE TO
5-METHYLTETRAHYDROFOLATE BY THE RAT JEJUNUM



EDWARD JAY OLINGER

1972

MUDD
LIBRARY
Medical

YALE



MEDICAL LIBRARY



Digitized by the Internet Archive
in 2017 with funding from
Arcadia Fund

<https://archive.org/details/conversionofpter00olin>

CONVERSION OF PTEROYLMONOGLUTAMATE TO 5-METHYLTETRAHYDROFOLATE

BY THE RAT JEJUNUM

by

Edward Jay Olinger

A dissertation presented to the faculty of the
Medical School of Yale University
in candidacy for the degree of Doctor of Medicine

New Haven, Connecticut

1972

ACKNOWLEDGMENTS

I wish first of all to thank Dr. Henry Binder and Dr. Joseph Bertino whose patience and counsel were the guiding light of this project. But, above all, it was their genuine friendship that made this entire effort valuable.

Grateful acknowledgment is made to Dr. Howard Spiro, for three years my friend and mentor.

I am also indebted to Mrs. Arlene Cashmore, Mrs. Judy Uhoch, Dr. Paul Chello and Claudia Rawlins for their expert assistance and good humor.

I wish to express gratitude to the American Society of Clinical Nutrition for their financial support of this work.

Finally, this acknowledgment would be incomplete without mentioning the support and encouragement of my wife.

TABLE OF CONTENTS

Summary	3
Introduction	5
Materials and Methods	8
Results	14
Discussion	16
Legends	21
Figures	23
Tables	32
References Cited	34

SUMMARY

Studies were designed to determine whether pteroylmono-glutamate (PGA) at physiologic concentrations is transported across the small intestine unaltered or is reduced and methylated to the circulating folate form during absorption. ^3H -PGA at varying concentrations was incubated on the mucosal side of rat jejunum placed between lucite chambers. Of the folate transferred to the serosal side, the percent identified as 5-methyltetrahydrofolate (5-MTHF) by DEAE-Sephadex chromatography was inversely related to the initial mucosal PGA concentration: at 7, 20 and 2000 nM, 46%, 32% and $<4\%$ respectively was 5-MTHF. In contrast, at 20 nM the ileum converted $<4\%$ to 5-MTHF. 10^{-6}M methotrexate, a dihydrofolate (DHF) reductase inhibitor, markedly inhibited PGA conversion to 5-MTHF by the jejunum. DHF reductase activity was measured in homogenates of villus scrapings to determine if the difference in PGA conversion to 5-MTHF between jejunum and ileum might be explained by differences in this enzyme activity. The purity of the villus preparation was confirmed by a 5 fold increase in sucrase specific activity (S.A.) and a 20 fold decrease in thymidine kinase S.A. compared to mucosal homogenates. A significant difference in DHF reductase S.A. in jejunal (0.105 ± 0.012 $\mu\text{mole/hr/mg}$ protein) compared to ileal villi (0.036 ± 0.003) was found. These studies demonstrate that both the reduction and methylation of PGA to 5-MTHF and DHF reductase S.A. is greater in the jejunum than in the ileum. The initial reduction

of PGA to tetrahydrofolate by DHF reductase may be the rate limiting step. These results could explain previous studies which have failed to identify conversion of PGA at higher and less physiologic concentrations.

INTRODUCTION

The absorption of pteroylglutamic acid (folic acid, PGA) in man after oral administration has been demonstrated by studies measuring plasma urine and fecal levels of folate derivatives¹⁻¹⁰. Clinical¹¹⁻¹⁵ and experimental¹⁶⁻²⁰ evidence favors the jejunum as the primary site of folic acid absorption although one study²¹ in the rat finds no difference between jejunal and ileal PGA absorption. However, the mechanism by which PGA and its derivatives are transported across the small intestine is still uncertain.

To understand the transport of folate*, it is useful to review qualitatively and quantitatively the distribution of folate derivatives in the diet and in the circulation. Analysis of food stuffs has revealed that between 80 and 90% of natural folates exist as polyglutamyl conjugates with unreduced and unconjugated PGA representing between 1 and 10% of the total dietary folate (500-700 µg)^{29,30}. Moreover, the predominant circulating and storage forms of folate are reduced forms, principally 5-methyltetrahydrofolate (5-MTHF)⁶⁵.

The data concerning the intestinal absorption of conjugated folate (figure 6) suggests that the polyglutamate is quantitatively deconjugated before release into the circulation^{9,31-35} although two studies^{36,37} give evidence for the

*refers to all forms of folate

absorption of the intact heptaglutamate. In addition, the finding of negligible conjugase activity in the intestinal brush border³⁸ has left the anatomical site of polyglutamate deconjugation in doubt. One study in rats²⁸ has elegantly demonstrated by a variety of techniques that for 5-MTHF, intestinal transport is by a non-energy dependent non-saturable system.

Despite our knowledge that 10% or less of dietary folate is PGA, most studies have dealt with the absorption of this compound alone^{16-21,23-27,45,46} since it is stable and available commercially. These investigations have presented conflicting data concerning the mechanism of PGA transfer across the intestine. Although several studies in rats^{16,21,23} and hamsters^{24,25} have demonstrated active transport as the process governing PGA absorption, other studies in the rat^{19,20,26} and hamster²⁷ have not demonstrated active transport and have suggested that PGA is absorbed by passive diffusion.

Following oral PGA administration, the plasma 5-MTHF level increases^{7,39,45,46}. Therefore PGA is converted to 5-MTHF at one or more sites between the point of ingestion and the peripheral circulation. Baker, et. al.⁷ suggested that folic acid was converted to 5-MTHF during transfer across the intestine, and studies in humans³⁹ and with everted intestinal sacs²² in the rat⁴⁰ and hamster²⁵ are in agreement with this thesis. In addition, the finding of DHF reductase activity in the small intestine⁴¹⁻⁴³ and its localization to the crypts of Lieberkuhn and the villus as determined by autoradiographic techniques⁴⁴

support those observations. However, in vivo perfusion studies of isolated intestinal segments in dogs³⁵, experiments with everted intestinal sacs in rats²⁰ and human studies^{45,46} involving hepatic and umbilical vein catheterization give evidence that PGA crosses the intestine intact without metabolic conversion.

The experiments failing to demonstrate reduction and methylation of PGA during absorption must be understood in light of the fact that conversion may have been masked by the pharmacologic doses of PGA used in these studies^{20,35,45,46} and that measures to protect reduced folates from light³⁵ and aerobic conditions^{20,35,45,46} appear not to have been adequately taken.

It is the purpose of this study, utilizing recently developed chromatographic methods and the apparatus described by Schultz and Zalusky⁴⁷, to determine whether PGA at physiologic concentrations is converted to other folate forms during transfer across the rat intestinal mucosa, and if so, to study some of the characteristics of this conversion.

MATERIALS AND METHODS

MATERIALS

Glucose oxidase, peroxidase, and o-dianisidine were purchased from Sigma, St. Louis, Missouri. $MgCl_2$ was purchased from Mallinckrodt, St. Louis, Missouri. ATP, creatine phosphokinase, creatine phosphate and unlabeled thymidine were purchased from Calbiochem, Los Angeles, California. Unlabeled folic acid (Folvite) and methotrexate (MTX) were obtained through the courtesy of Lederle, Pearl River, New York. 2,5-Diphenyloxazole, p-Bis[2-(5-phenyloxazolyl)]-benzene and thymidine-2- ^{14}C (54 $\mu Ci/\mu mole$) were purchased from New England Nuclear, Boston, Massachusetts. 3H -PGA (33 Ci/mmole and 29 Ci/mmole) was purchased from Amersham-Searle, Des Plaines, Illinois.

METHODS

I. Determination of mucosal to serosal folate transfer in the rat small intestine.

The two lots of 3H -PGA labeled in the 3', 5' and 9 positions of the molecule (figure 6) were purified by column chromatography on A-25 DEAE-Sephadex (Pharmacia). The purified lots had specific activities of 5.6 Ci/mmole and 45.8 Ci/mmole respectively, and were greater than 99% radiochemically pure.

Mucosal to serosal transfer of 3H -PGA and its derivatives were determined under short circuited conditions using the methods and apparatus described by Schultz and Zalusky⁴⁷. Flat sheets of rat jejunum or ileum from male albino Sprague-Dawley

rats weighing 300-350 grams were mounted between lucite chambers. Equimolar amounts of unlabeled PGA at varying concentrations were placed on both the mucosal and serosal side of the chamber with approximately 3.0 μ Ci of ^3H -PGA added to the mucosal solution. After a 20 minute equilibration period, the rate of appearance of radiolabeled material in the serosal solution was determined by taking 1 ml aliquots from the serosal media at 20 minute intervals. The sample added to 10 ml of Bray's solution⁴⁸ was counted in a Packard Tri-Carb 3-Channel liquid scintillation spectrometer for 10 minutes. In these studies, results were expressed in pmoles of folate transferred to the serosal solution per cm^2 intestine per 100 minutes. All experiments were performed in the absence of light. Both sides of the tissue were bathed with identical solutions maintained at 37°C . and bubbled continuously with $\text{O}_2\text{-CO}_2$ (95:5, v/v). In all experiments, the composition of the buffer bathing solution in millimoles per liter was sodium chloride, 115; sodium bicarbonate, 25; potassium phosphate, 5; magnesium chloride, 1.2; calcium chloride, 1.2; ascorbic acid, 17; pH 7.4.

II. Identification of folate compounds by chromatography.

At the end of 100 minutes, the serosal bathing solutions from two to three experiments were pooled and stored overnight in 3 M mercaptoethanol at 4°C . A similar procedure was followed for the mucosal bathing solutions. Folate compounds were chromatographically identified using the method described by Nixon and Bertino⁴⁹. The pooled serosal or mucosal samples containing folate markers were diluted to a salt concentration of 0.1 molar

and applied to a 0.9 x 27 cm column of A-25 DEAE-Sephadex equilibrated with 0.1 M phosphate buffer, pH 6.0, containing 10 mM mercaptoethanol. Materials were eluted by phosphate buffer containing 10 mM mercaptoethanol, pH 6.0, the concentration of which was increased linearly from 0.1 M to 1.8 M. The final concentration of 1.8 M phosphate buffer was obtained after passage of 30 ml of eluting buffer. Non-radioactive markers of 5-MTHF, PGA and p-aminobenzoylglutamate (pABG) were added with the sample. Fractions of 2.5 ml were collected in the dark at 4°C. and monitored by ultraviolet absorbance spectra and by radioactivity. 0.5 ml aliquots per fraction were counted with the Packard Tri-Carb spectrometer in 15 ml of scintillation fluid containing 8.0 g of 2,5-Diphenyloxazole in 1 liter of absolute alcohol and 100 mg of p-Bis[2-(5-phenyloxazolyl)]-benzene in 2 liters of toluene. The compounds identified were quantitated in terms of pmoles of PGA or 5-MTHF transferred to the serosal solution per cm² intestine per 100 min by integrating the area under each peak in the chromatograph to determine the relative proportion of each compound transferred, and by combining this information with the total mucosal to serosal folate transfer obtained as described in section I. The average number of counts added to the column to identify the various folate compounds was between 25,000-30,000 cpm. At this level of activity, a 5-MTHF peak of at least 4% of the total folate activity can be identified. Therefore, in those experiments in which a 5-MTHF peak was not observed, we are assuming that less than 4% 5-MTHF

was present. Obviously, the correct amount of 5-MTHF is somewhere between 0 and 4%.

III. Preparation of rat small intestine for enzyme analysis and histology.

Male albino Sprague-Dawley rats weighing 200-250 g placed under ether anesthesia had their small bowel excised and placed in cold $0.5 \times 10^{-4}M$ tris buffer with $10^{-4}M$ thymidine*, pH 7.5 (tris-thymidine buffer). The small intestine was divided into three equal parts of about 30 cm in length. The intestinal contents were removed from the proximal (jejunal) and distal (ileal) segments by flushing with cold tris-thymidine buffer and the middle segment was discarded. The jejunum was divided into three equal parts of about 10 cm in length and two of the three segments selected at random were lightly scraped along the mucosa according to the method of Dietchy and Siperstein⁵⁰ to separate the intestinal villi from the crypts-muscle layer. Jejunal villi scrapings were pooled and placed in 2-3 ml of tris-thymidine buffer. The third 10 cm segment underwent a more forceful scraping to separate total mucosa from the underlying

*In preliminary experiments, it was found that the thymidine kinase assay could be performed on the intestinal homogenates only if the enzyme was stabilized by the presence of $10^{-4}M$ thymidine in tris buffer and if fresh supernatant fractions were utilized for the assay.

smooth muscle layer, after which the total mucosa was placed in 3-5 ml of tris-thymidine buffer. The same procedures were followed for the 30 cm ileal segment. The histology of these intestinal layers was determined on formalin fixed, hematoxylin and eosin and masson stained sections of each preparation (figure 9). The jejunal and ileal villi and total mucosa were each individually homogenized for 1-3 minutes with a glass motor-driven pestle in the initial volume of tris-thymidine buffer. The homogenates were then centrifuged for 15 minutes at 27,000 x g and the supernatant from each homogenate was divided into three aliquots. One aliquot was saved for the immediate determination of thymidine kinase activity and the other two were stored at -70°C. until used (usually within three days), for the determination of dihydrofolate reductase activity, sucrase activity and protein.

Thymidine kinase assay: The thymidine kinase activity of the fresh supernatant fraction of villi and total mucosa homogenates was determined using the assay of Chello and Jaffe⁵¹ and the chromatographic separation of Furlong⁵². The thymidine kinase activity is expressed as pmoles of thymidine-5'-monophosphate formed per mg protein per 30 minutes.

Dihydrofolate reductase assay: The dihydrofolate reductase activity of the supernatant fractions of villi and total mucosa homogenates was determined at pH 7.5 at 37°C. using the method of Bertino, et. al.⁵³. Activity is expressed as pmoles of dihydrofolate reduced per hour per mg protein.

Sucrase assay: The sucrase activity of the supernatant frac-

tions of villi and total mucosa homogenates was determined according to the method of Dahlqvist⁵⁴. Activity is expressed as μ g of glucose liberated per hour per mg protein.

Protein determination: Protein concentrations of the supernatant fractions of villi and total mucosa homogenates were determined by the biuret method with bovine serum albumin as the standard.

Statistics: Statistical significance was determined by the student t test⁵⁵.

RESULTS

The identification and quantitation of folate compounds absorbed by rat small intestine.

The DEAE-Sephadex chromatographic identification of folate compounds in the serosal bathing solutions of rat jejunum incubated with 7 nM PGA, 20 nM PGA, 20nM PGA with 10^{-6} M MTX, and with 2000 nM PGA and the ileum with 20 nM PGA are shown in figures 1-5. The folate transfer studies were combined with this chromatographic data to give the pmoles of 5-MTHF and PGA transferred per cm^2 intestine per 100 min as shown in table 1. The total folate transfer was found to be linearly related to mucosal PGA concentration (table 1 and figure 8).

It is seen that when the concentration of PGA in the mucosal bathing solution incubating the jejunum is increased from 7 nM to 2000 nM, the per cent of the total folate identified in the serosal bathing solution as 5-MTHF decreased from 46% to less than 4%. Further, the presence of 10^{-6} M MTX, a dihydro-folate reductase inhibitor, markedly decreased the appearance of 5-MTHF in the serosal solution when 20 nM PGA was present on the mucosal side of the jejunum. When 20 nM PGA was on the mucosal side of the ileum, less than 4% 5-MTHF was identified.

The pABG peak identified for the serosal media is the result of oxidative breakdown of PGA and reduced folates occurring under the incubating conditions employed in these experiments (figure 7). The possibility of selective pABG uptake by intestinal mucosa awaits further study.

Enzyme analysis of small intestine.

The thymidine kinase, DHF reductase and sucrase specific activities of the supernatant fractions of jejunal and ileal villi and total mucosa are shown in table 2. The purity of the villus preparation was confirmed by a five fold increase in sucrase specific activity and a twenty fold decrease in thymidine kinase specific activity when compared to total mucosa homogenates and by histology (figure 9). Although DHF reductase specific activity was similar in proximal and distal total mucosa homogenates ($P < 0.2$), a three fold increase in DHF reductase specific activity was found in jejunal villi compared to ileal villi ($P < 0.001$). On the other hand, the jejunal:ileal ratio of thymidine kinase and sucrase specific activities was identical for both total mucosa homogenates and villi.

DISCUSSION

These studies demonstrate that the reduction and methylation of PGA to 5-MTHF occur in the rat small intestine and that this conversion is significantly greater in the jejunum than in the ileum. It is noted that in no chromatogram (figures 1-5) were peaks of radioactivity identified which might prove to be intermediate compounds in the pathway of PGA's conversion to 5-MTHF. This suggests that the initial reduction of PGA to tetrahydrofolate (FH_4) by DHF reductase may be the rate limiting step in this pathway (figure 7). However the absence of these peaks may indicate that the 17 mM ascorbate in the serosal bathing solution was not in sufficiently high concentration to protect these intermediate compounds from oxidation, and/or that chromatographing these intermediates in amounts smaller than the assay system was capable of identifying may have resulted in their presence being masked.

That methotrexate, a competitive stoichiometric inhibitor⁵⁶ of DHF reductase, markedly diminished the percent PGA converted to 5-MTHF by the rat jejunum (figure 3) suggested to us that the differences in PGA reduction and methylation to 5-MTHF between jejunum and ileum might be explained by differences in DHF reductase activity in these two regions of small intestine. In experiments with supernatant fractions of total mucosa homogenates from jejunum and ileum, the differences in DHF reductase activity was not statistically significant ($P < 0.2$). However, the crypt region within the mucosa of the small bowel, a region

of active DNA^{57,58} and protein^{57,59} synthesis, might be expected to contain this enzyme in sufficient activity to mask real differences in specific activity between jejunal and ileal villi, the tissue usually considered important in intestinal absorption. Villus homogenates prepared according to the method of Dietchy and Siperstein⁵⁰ did reveal a higher DHF reductase specific activity in jejunal compared to ileal villi ($P < 0.001$). This is in contrast to findings of an identical jejunal:ileal ratio for thymidine kinase and sucrase specific activities in both total mucosa homogenates and villi (table 2).

The use of sucrase and thymidine kinase as enzyme markers to verify the purity of the villus preparation is based on the work of Van Genderen⁶⁰ and others^{57,61,62}, which dealt with sequential biochemical analyses from villus to crypt in the intestine of the rat and demonstrated villi:crypt ratios of 4.0 and 0.02 for sucrase⁵⁷ and thymidine kinase⁶² respectively.

There is at least one study which by autoradiographic techniques⁴⁴ indirectly demonstrated DHF reductase activity in the villi of the proximal small bowel. Results of further experiments to determine if DHF reductase is localized to the microvilli of the brush border might suggest, as proposed by Crane⁶³, whether or not a "kinetic advantage" exists for the absorption of PGA similar to that which may exist for disaccharide absorption, on the basis of specific membrane-substrate-enzyme organization at the level of the microvillus. It would also be interesting to determine whether DHF reductase in the highly specialized villus has the same properties as DHF reductase in the relatively

undifferentiated crypt cell.

These results indicate that the transfer of PGA across the intestine is directly related to the mucosal PGA concentration (figure 8). However the percent PGA converted to 5-MTHF is not linearly related to PGA concentration: at 7 nM PGA, 46% is methylated, but at 2000 nM less than 4% can be identified as 5-MTHF in the serosal solution (table 1). This data can then be interpreted to indicate that PGA to 5-MTHF conversion requires a rate limiting step but that total PGA transfer does not take place in a saturable system (figure 8). One would like to speculate that the saturable process for PGA absorption observed by some investigators^{16,20,21,23-25} is related to this rate limiting step of PGA conversion to 5-MTHF.

Therefore, what one finds in the serosal solution or pre-hepatic blood sample is dependent upon the concentration of PGA used in the mucosal incubating media, the surface area and time available for absorption, the quantity of folate in the sample to be analyzed and upon the sensitivity of the assay method employed. Previous experiments in the field which appeared to be utilizing quantities of PGA greater than 100 nM^{20,35,45,46} would therefore not be expected to identify a significant percentage of 5-MTHF during absorption. Smith, Matty and Blair, in studies with everted intestinal sacs from rats²⁰ incubated with 10^{-7} to 10^{-6} M PGA, did report a "minor radioactive component" with their chromatographic assay system which cochromatographed with both 5-formyltetrahydropteroylglutamic acid and 5-MTHF markers. Also, the possibility that the measures taken in

these studies to protect reduced folates from light³⁵ and aerobic conditions^{20,35,45,46} were not optimal may have contributed to the difficulty encountered by these workers in identifying absorbed 5-MTHF. Both Strum, et. al.⁴⁰ and Cohen²⁵, having used everted intestinal sacs from rats and hamsters respectively, found conversion of PGA to 5-MTHF during absorption. However, they did not study the characteristics of this conversion.

The mucosa to serosa transfer data for jejunum and ileum (table 1), derived from only two to three pooled experiments per study, does not have statistical validity for any conclusion to be drawn regarding a preferential site for folate absorption. Past clinical¹¹⁻¹⁵ and experimental¹⁶⁻²⁰ research favors the jejunum as the primary site of folic acid absorption although one study in the rat²¹ finds no difference between jejunal and ileal PGA absorption.

The recent observation⁶⁴ that the small intestine of the rat is the only organ which lacks the 5-methyltetrahydrofolate-homocysteine methyltransferase enzyme for the conversion of homocysteine to methionine may indicate that in methionine metabolism there is a shunting of 5-MTHF, the predominant circulating and storage form of folate⁶⁵, via the blood stream to the liver, kidney, adrenals, testes, pancreas and brain, where in all these organs good specific activity for this enzyme has been found⁶⁴. If other enzymes related to folate utilization are found to be lacking in the small intestine, it may suggest that the small intestine is an effective "scavenger" of dietary folate for preferential use elsewhere in the body.

Reports of a congenital isolated defect for folate absorption in man^{66,67} offer a good opportunity for further elucidation of the specific uptake mechanisms for PGA by the mucosal cell.

LEGENDS

- Figure 1. Chromatographic identification on DEAE-Sephadex of folate compounds found in the serosal bathing solution of rat jejunum mounted between lucite chambers and incubated with 7 nM PGA in the mucosal and serosal solutions with ^3H -PGA in the mucosal solution. Results of three pooled experiments. Radioactivity was identified in the 273, 280 and 289 m μ peaks which cochromatographs with unlabeled pABG, PGA and 5-MTHF. pABG, PGA and 5-MTHF were added to the column as markers.
- Figure 2. Chromatographic identification on DEAE-Sephadex of folate compounds found in the serosal bathing solution of rat jejunum incubated with 20 nM PGA. Results of two pooled experiments. Further details are presented in figure 1 and in the text.
- Figure 3. Chromatographic identification on DEAE-Sephadex of folate compounds found in the serosal bathing solution of rat jejunum incubated with 20 nM PGA and 10^{-6}M methotrexate. Results of two pooled experiments. Further details are presented in figure 1 and in the text.
- Figure 4. Chromatographic identification on DEAE-Sephadex of folate compounds found in the serosal bathing solution of rat jejunum incubated with 2000 nM PGA. Results of two pooled experiments. Further details

are presented in figure 1 and in the text.

Figure 5. Chromatographic identification on DEAE-Sephadex of folate compounds found in the serosal bathing solution of rat ileum incubated with 20 nM PGA. Results of two pooled experiments. Further details are presented in figure 1 and in the text.

Figure 6. The structure of pterioic acid, folic acid and folic acid polyglutamates. Glutamic acid residues are gamma carboxy linked. Modified from Krumdieck and Baugh⁶⁸.

Figure 7. The pathway for the reduction and methylation of folic acid to 5-methyltetrahydrofolic acid.

Figure 8. Graphic representation of data from table 1 demonstrating the transfer of total folate (—) and 5-MTHF (----) from the mucosal to the serosal solution with increasing $[PGA]_{\text{mucosal}}$.

Figure 9. Histologic demonstration of villus scrapings prepared as described in the text. Only pure villi were identified in the villus scrapings.

figure 1.




figure 2.




figure 3.




figure 4.




figure 5.




FIGURE 6.




FIGURE 7.




figure 8.




Figure 9.



14	1000	1000
15	1000	1000
16	1000	1000
17	1000	1000
18	1000	1000
19	1000	1000
20	1000	1000
21	1000	1000
22	1000	1000
23	1000	1000
24	1000	1000
25	1000	1000
26	1000	1000
27	1000	1000
28	1000	1000
29	1000	1000
30	1000	1000
31	1000	1000
32	1000	1000
33	1000	1000
34	1000	1000
35	1000	1000
36	1000	1000
37	1000	1000
38	1000	1000
39	1000	1000
40	1000	1000
41	1000	1000
42	1000	1000
43	1000	1000
44	1000	1000
45	1000	1000
46	1000	1000
47	1000	1000
48	1000	1000
49	1000	1000
50	1000	1000
51	1000	1000
52	1000	1000
53	1000	1000
54	1000	1000
55	1000	1000
56	1000	1000
57	1000	1000
58	1000	1000
59	1000	1000
60	1000	1000
61	1000	1000
62	1000	1000
63	1000	1000
64	1000	1000
65	1000	1000
66	1000	1000
67	1000	1000
68	1000	1000
69	1000	1000
70	1000	1000
71	1000	1000
72	1000	1000
73	1000	1000
74	1000	1000
75	1000	1000
76	1000	1000
77	1000	1000
78	1000	1000
79	1000	1000
80	1000	1000
81	1000	1000
82	1000	1000
83	1000	1000
84	1000	1000
85	1000	1000
86	1000	1000
87	1000	1000
88	1000	1000
89	1000	1000
90	1000	1000
91	1000	1000
92	1000	1000
93	1000	1000
94	1000	1000
95	1000	1000
96	1000	1000
97	1000	1000
98	1000	1000
99	1000	1000
100	1000	1000

TABLE 1.

Jejunal Conversion of PGA to 5-MTHF

Study #	Tissue	nM PGA in mucosal solution	Total** Folate J* M \rightarrow S (pm/cm ² /100 min)	% 5-MTHF of total J Folate M \rightarrow S	J 5-MTHF M \rightarrow S (pm/cm ² /100 min)	% PGA of total J Folate M \rightarrow S	J PGA M \rightarrow S (pm/cm ² /100 min)
1	Jejunum	7	0.103	46	0.047	54	0.056
2	Jejunum	20	0.260	34	0.090	66	0.170
3	Jejunum	20 plus 10 ⁻⁶ M MTX	0.424	6	0.026	94	0.398
4	Jejunum	2000	47.738	< 4	< 1.910	> 96	> 45.828
5	Ileum	20	0.557	< 4	< 0.022	> 96	> 0.535

** PGA and 5-MTHF

* Transfer from mucosal to serosal solution

4
5
6
7
8

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

10

TS:

1

200

13

TABLE 2.

DHF reductase, sucrase and thymidine kinase specific activities (S.A.) in the supernatant fractions of villi and total mucosa homogenates from rat jejunum and ileum (\pm S.E.)

Tissue (n=6)	DHF reductase S.A.		Sucrase S.A.		Thymidine kinase S.A.	
	(μ moles DHF reduced/ mg protein/hr at 37°C)		(μ g glucose liberated/mg protein/hr)		(pmoles TMP formed/mg protein/30 min)	
jejunal villi	0.105* \pm 0.012		68.20 \pm 3.66		4.31 \pm 0.91	
jejunal total mucosa	0.235** \pm 0.040		14.43 \pm 2.77		115.90 \pm 24.96	
ileal villi	0.036 \pm 0.003		67.82 \pm 5.34		4.03 \pm 0.89	
ileal total mucosa	0.155 \pm 0.030		14.87 \pm 1.39		102.59 \pm 11.74	

** p < 0.2, compared to ileal total mucosa

* p < 0.001, compared to ileal villi

Mr. J. Edgar Hoover, Director
Federal Bureau of Investigation, Washington, D.C.

Dear Mr. Hoover:

I am writing to you regarding the matter of the
recently reported disappearance of the
body of the late Mr. J. Edgar Hoover.

I am sure that you will be interested in the
fact that the body of the late Mr. J. Edgar Hoover
has been found.

I am sure that you will be interested in the
fact that the body of the late Mr. J. Edgar Hoover
has been found.

Sincerely,
J. Edgar Hoover

REFERENCES CITED

1. Denko, C.W. 1951. Pteroylglutamic Acid Clearance in Normal Adults. *J. Appl. Physiol.* 3:559.
2. Spray, G.H. and Witts, L.J. 1952. The Utilization of Folic Acid Given by Mouth. *Clin. Sci.* 11:273.
3. Clark, S.L., Jr. 1953. Oral Folic Acid Tolerance Test in Normal Human Subjects and Patients with Pernicious Anemia. *Proc. Soc. Exp. Biol. (N.Y.)* 82:25.
4. Butterworth, L.E., Jr., Nadel, H., Perez-Santiago, E., Santini, R., Jr., and Gardner, F.H. 1957. Folic Acid Absorption, Excretion and Leukocyte Concentrations in Tropical Sprue. *J. Lab. Clin. Med.* 50:673.
5. Chanarin, I., Anderson, B.B. and Mollin, D.L. 1958. The Absorption of Folic Acid. *Brit. J. Haemat.* 4:156.
6. Girdwood, R.H. 1960. Folic Acid, its Analogs and Antagonists. *Advanc. Clin. Chem.* 3:235.
7. Baker, H., Frank, O., Feingold, S., Ziffer, H., Gellene, R.A., Leevy, C.M. and Sobotka, H. 1965. The Fate of Orally and Parenterally Administered Folates. *Amer. J. Clin. Nut.* 17:88.
8. Steinkamp, R., Shukers, C.F., Totter, J.R. and Day, P.L. 1946. Urinary Excretion of Orally Administered Pteroylglutamic Acid. *Proc. Soc. Exptl. Biol. Med.* 63:556.
9. Swendsøid, M.E., Bird, O.D., Brown, R.A. and Bethell, F.H. 1947. Metabolic Function of Pteroylglutamic Acid and its Hexaglutamyl Conjugate. II. Urinary Excretion Studies on Normal Persons. Effect of a Conjugase Inhibitor. *J. Lab. Clin. Med.* 32:23.
10. Jukes, T.H., Franklin, A.L., Stokstad, E.L.R. and Boehne, J.W. 1947. The Urinary Excretion of Pteroylglutamic Acid and Certain Related Compounds. *J. Lab. Clin. Med.* 32:1350.
11. Klipstein, F.A. 1966. Folate Deficiency Secondary to Disease of the Intestinal Tract. *Bull. N.Y. Acad. Med.* 42:638.
12. Cox, E.V., Meynell, M.J. and Cooke, W.T. 1958. The Folic Acid Excretion Test in the Steatorrhea Syndrome. *Gastroenterology* 35:390.

13. Klipstein, F.A. 1963. The Urinary Excretion of Orally Administered Tritium-Labeled Folic Acid as a Test of Folic Acid Absorption. *Blood* 21:626.
14. Baker, H., Thomson, A., Feingold, S. et. al. 1969. Role of the Jejunum in the Absorption of Folic Acid and its Polyglutamates. *Amer. J. Clin. Nut.* 22:124.
15. Kinnear, D.G., Johns, D.G., MacIntosh, P.C. et. al. 1963. Intestinal Absorption of Tritium-Labeled Folic Acid in Idiopathic Steatorrhea: Effect of a Gluten-Free Diet. *Canad. Med. Asso. J.* 89:975.
16. Herbert, V. and Shapiro, S.S. 1962. The Site of Absorption of Folic Acid in the Rat In Vitro. *Fer. Proc.* 21:260.
17. Hepner, G.W. 1969. The Absorption of Pteroylglutamic (Folic) Acid in Rats. *Brit. J. Haemat.* 16:241.
18. Hepner, G.W., Booth, C.C., Cowan, J. et. al. 1968. Absorption of Crystalline Folic Acid in Man. *Lancet* 2:302.
19. Yoshino, T. 1968. The Clinical and Experimental Studies on the Metabolism of Folic Acid Using Tritiated Folic Acid. II. The Experimental Studies on the Absorption Site and Mechanism of Tritiated Folic Acid in Rats. *J. Vitam.* 14:35.
20. Smith, M.E., Matty, A.J. and Blair, J.A. 1970. The Transport of Pteroylglutamic Acid Across the Small Intestine of the Rat. *Biochim. Biophys. Acta.* 219:37.
21. Burgen, A.S.V. and Goldberg, N.J. 1962. Absorption of Folic Acid from the Small Intestine of the Rat. *Brit. J. Pharmacol.* 19:313.
22. Wilson, T.H. and Wiseman, G. 1954. The Use of Sacs of Everted Small Intestine for the Study of the Transference of Substances from the Mucosal to the Serosal Surface. *J. Physiol.* 123:116.
23. Herbert, V. 1967. Biochemical and Hematologic Lesions in Folic Acid Deficiency. *Am. J. Clin. Nut.* 20:562.
24. Cohen, N., Gelb, A. and Sobotka, H. 1964. Intestinal Absorption of Folic Acid In Vitro (Abstr.). *Clin. Res.* 12:206.
25. Cohen, N. 1965. Differential Microbiological Assay in Study of Folic Acid Absorption In Vitro by Everted Intestinal Sacs (Abstr.). *Clin. Res.* 13:252.

1. The first part of the document is a list of names and titles, including "The Hon. Mr. Justice" and "The Hon. Mr. Justice".

26. Turner, J.B. and Hughs, D.E. 1962. The Absorption of Some B-Group Vitamins by Surviving Rat Intestine Preparations. Quart. J. Exptl. Physiol. 47:107.
27. Spencer, R.P. and Bow, T.M. 1964. In Vitro Transport of Radiolabeled Vitamins in the Small Intestine. J. Nuclear Med. 5:251.
28. Strum, W., Nixon, P., Bertino, J. and Binder, H. 1971. Intestinal Folate Absorption. I. 5-Methyltetrahydrofolic Acid. J. Clin. Invest. 50:1910.
29. Butterworth, C.E., Jr., Santini, R. Jr., and Frommeyer, W. B., Jr. 1963. The Pteroylglutamate Components of American Diets as Determined by Chromatographic Fractionation. J. Clin. Invest. 42:1929.
30. Perry, J. 1971. Folate Analogues in Normal Mixed Diets. Brit. J. Haemat. 21:435.
31. Bethell, F.H., Meyers, M.C., Andrews, G.A. et. al. 1947. Metabolic Function of Pteroylglutamic Acid and its Hexa-glutamyl Conjugate. I. Hematologic and Urinary Excretion Studies in Patients with Macrocytic Anemia. J. Lab. Clin. Med. 32:3.
32. Perry, J. and Chanarin, I. 1968. Absorption and Utilization of Polyglutamyl Forms of Folate in Man. Brit. Med. J. 4:546.
33. Butterworth, C.E., Jr., Baugh, C.M. and Krumdieck, C. 1969. A Study of Folate Absorption and Metabolism in Man Utilizing Carbon-14 labeled Polyglutamates Synthesized by the Solid Phase Method. J. Clin. Invest. 48:1131.
34. Bernstein, L.H., Gutstein, S., Weiner, S. et. al. 1970. The Absorption and Malabsorption of Folic Acid and its Polyglutamates. Amer. J. Med. 48:570.
35. Baugh, C.M., Krumdieck, C.L., Baker, H.J. and Butterworth, C.E., Jr. 1971. Studies on the Absorption and Metabolism of Folic Acid. I. Folate Absorption in the Dog after Exposure of Isolated Intestinal Segments to Synthetic Pteroyl-polyglutamates of Various Chain Lengths. J. Clin. Invest. 50:2009.
36. Cooperman, J.M. and Lubby, A.L. 1965. The Physiological Fate in Man of Some Naturally Occurring Polyglutamates of Folic Acid. Israel J. Med. Sci. 1:704.

Mr. J. C. ... and ...
...

... ..
... ..
... ..

... ..
... ..

... ..
... ..
... ..
... ..

37. Benerjee, D.K. and Chatterjea, J.B. 1966. Observations on the Presence of Pteroylpolyglutamates in Human Serum. Blood 28:913.
38. Hoffbrand, A.V. and Peters, T.J. 1969. The Subcellular Localization of Pteroylpolyglutamate Hydrolase and Folate in Guinea Pig Intestinal Mucosa. Biochim. Biophys. Acta. 192:479.
39. Perry, J. and Chanarin, I. 1970. Intestinal Absorption of Reduced Folate Compounds in Man. Brit. J. Haemat. 18:329.
40. Strum, W.B., Nixon, P.F., Binder, H.J. et. al. 1970. Intestinal Absorption of 5-Methyltetrahydrofolate. (Abstr.). Clin. Res. 18:389.
41. Werkheiser, W.C. 1961. Specific Binding of 4-Amino Folic Acid Analogues by Folic Acid Reductase. J. Biol. Chem. 236:888.
42. Bertino, J.R., Simmons, B. and Donohue, D.M. 1964. Levels of Dehydrofolate Reductase and the Formate-Activating Enzyme Activities in Guinea Pig Tissues before and after Amethopterin Administration. Biochem. Pharm. 13:225.
43. Bertino, J.R., Iannotti, A.T., Perkins, J.P. and Johns, D.G. 1966. Dihydrofolate Reductase from Guinea Pig Liver and Small Intestine. Biochem. Pharm. 15:563.
44. Darzynkiewicz, Z. Rogers, A.W., Barnard, E.A., Wang, D.-H. and Werkheiser, W.C. 1966. Autoradiography with Tritiated Methotrexate and the Cellular Distribution of Folate Reductase. Science 151:1528.
45. Whitehead, V.M. and Cooper, B.A. 1967. Absorption of Unaltered Folic Acid from the Gastro-Intestinal Tract in Man. Brit. J. Haemat. 13:679.
46. Melikian, V., Paton, A., Leeming, R.J. and Portman-Graham, H. 1971. Site of Reduction and Methylation of Folic Acid. Lancet 2:955.
47. Schultz, S.G. and Zalusky, R. 1964. Ion Transport in Isolated Rabbit Ileum. I. Short-Circuit Current and Na Fluxes. J. Gen. Physiol. 47:567.
48. Bray, G.A. 1960. A Simple Efficient Liquid Scintillator for Counting Aqueous Solutions in a Liquid Scintillation Counter. Anal. Biochem. 1:279.

1000 1000 1000 1000 1000 1000 1000 1000 1000 1000

1000 1000 1000 1000 1000 1000 1000 1000 1000 1000

1000 1000 1000 1000 1000 1000 1000 1000 1000 1000

1000 1000 1000 1000 1000 1000 1000 1000 1000 1000

1000 1000 1000 1000 1000 1000 1000 1000 1000 1000

1000 1000 1000 1000 1000 1000 1000 1000 1000 1000

49. Nixon, P.F. and Bertino, J.R. 1971. Separation and Identification of Folate Co-Enzymes on DEAE-Sephadex. In Vitamins and Co-Enzymes, a Volume of Methods in Enzymology. D.B. McCormick and L.D. Wright, Editors. Acad. Press. New York.
50. Dietchy, J.M. and Siperstein, M.D. 1965. Cholesterol Synthesis by the Gastrointestinal Tract: Localization and Mechanism of Control. J. Clin. Invest. 44:1311.
51. Chello, P.L. and Jaffe, J.J. 1972. Isolation, Partial Purification and Properties of Thymidine Kinase from Trypanosoma (Trypanozoon) Brucci Rhodesiense. J. Parasitology (in press).
52. Furlong, M.B. 1963. A Rapid Assay for Nucleotide Kinases Using ¹⁴C or ³H Nucleotides. Anal. Biochem. 5:515.
53. Bertino, J.R., Perkins, J.P. and Johns, D.G. 1965. Purification and Properties of Dihydrofolate Reductase from Ehrlich Ascites Carcinoma Cells. Biochem. 4:839.
54. Dahlqvist, A. 1968. Assay of Intestinal Disaccharidases. Anal. Biochem. 22:99.
55. Snedecor, G.W. 1956. Statistical Methods Applied to Experiments in Agriculture and Biology. Iowa State University Press, Iowa. 5th Edition.
56. Blakley, R.L. 1969. The Biochemistry of Folic Acid and Related Pteridines. John Wiley and Sons Inc., New York. P. 158.
57. Das, B.C. and Gray, G.M. 1969. Protein Synthesis in Small Intestine: Localization and Correlation with DNA Synthesis and Sucrase Activity. Biochim. Biophys. Acta. 195:255.
58. Leblond, C.P. and Stevens, C.E. 1948. The Constant Renewal of the Intestinal Epithelium of the Albino Rat. Anat. Record. 100:357.
59. Lipkin, M. and Quastler, H. 1962. Studies of Protein Metabolism in Intestinal Epithelial Cells. J. Clin. Invest. 41:646.
60. Van Genderen, H. and Engels, C. 1938. On the Distribution of Some Enzymes in the Duodenum and Ileum of the Cat. Enzymologica 5:71.

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
530 SOUTH EAST ASIAN AVENUE

CHICAGO, ILLINOIS 60607
U.S.A.

PROFESSOR DR. J. H. DUNN
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF CHICAGO
530 SOUTH EAST ASIAN AVENUE
CHICAGO, ILLINOIS 60607

DR. J. H. DUNN
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF CHICAGO
530 SOUTH EAST ASIAN AVENUE
CHICAGO, ILLINOIS 60607

DR. J. H. DUNN
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF CHICAGO
530 SOUTH EAST ASIAN AVENUE
CHICAGO, ILLINOIS 60607

61. Nordstrom, C., Dahlqvist, A. and Josefsson, L. 1968. Quantitative Determination of Enzymes in Different Parts of the Villi and Crypts of Rat Small Intestine. Comparison of Alkaline Phosphatase, Disaccharidases and Dipeptidases. J. Histochem. Cytochem. 15:713.
62. Fortin-Magna, R., Hurwitz, R. Herbst, J.J. and Kretchmer, N. 1970. Intestinal Enzymes: Indicators of Proliferation and Differentiation in the Jejunum. Science 167:1627.
63. Crane, R.K. 1969. A Perspective of Digestive-Absorption Function. Amer. J. Clin. Nut. 22:242.
64. Finkelstein, J.D., Kyle, W.E. and Harris, B.J. 1971. Methionine Metabolism in Mammals. Regulation of Homocysteine Methyltransferase in Rat Tissue. Arch. Biochem. and Biophys. 146:84.
65. Noronha, J.M. and Silverman, M. 1962. Distribution of Folic Acid Derivatives in Natural Materials. I. Chicken Liver Folates. J. Biol. Chem. 237:3299.
66. Luhby, A.L., Eagle, F.J., Roth, E. et. al. 1961. Relapsing Megaloblastic Anemia in an Infant due to a Specific Defect in Gastrointestinal Absorption of Folic Acid. Amer. J. Dis. Child. 102:482.
67. Lanzkowsky, P., Erlandson, M.E. and Bezan, A.I. 1969. Isolated Defect in Folic Acid Absorption Associated with Mental Retardation and Cerebral Calcification. Blood 34:452.
68. Krumdieck, C.L. and Baugh, C.M. 1969. The Solid Phase Synthesis of Polyglutamates of Folic Acid. Biochem. 8:1568.

YALE MEDICAL LIBRARY

Manuscript Theses

Unpublished theses submitted for the Master's and Doctor's degrees and deposited in the Yale Medical Library are to be used only with due regard to the rights of the authors. Bibliographical references may be noted, but passages must not be copied without permission of the authors, and without proper credit being given in subsequent written or published work.

This thesis by _____ has been
used by the following persons, whose signatures attest their acceptance of the
above restrictions.

NAME AND ADDRESS

DATE

